

-

Xydous, M., Sekeri-Pataryas, K.E., Prombona, A. and T.G. Sourlingas. Nicotinamide treatment reduces the levels of histone H3K4 trimethylation in the promoter of the *mper1* circadian clock gene and blocks the ability of dexamethasone to induce the acute response. *BBA-Gene Reg. Mech.* 1819: 877-884, 2012.

-

Salpea, P., Russanova, V.R., Hirai, T.H., Sourlingas, T.G., Sekeri-Pataryas, K.E., Romero, R., Epstein, J. and B.H. Howard. Postnatal development- and age-related changes in DNA-methylation patterns in the human genome. *Nucleic Acids Res.* 40 (14): 6477-6494, 2012.

-

Repouskou, A, Sourlingas, T.G., Sekeri-Pataryas, K.E. and A. Prombona. The circadian expression of c-Myc is modulated by the histone deacetylase inhibitor trichostatin A in synchronized murine neuroblastoma cells. *Chronobiology Int.*, 27(4): 722-741, 2010.

-

Ninios, Y.P., Sekeri-Pataryas, K.E. and T.G. Sourlingas. Histone H1 subtype preferences of DFF40 and possible nuclear localization of DFF40/45 in normal and trichostatin A-treated NB4 leukemic cells. *Apoptosis*, 15(2): 128-138, 2010.

-

Ninios, Y.P., Sekeri-Pataryas, K.E. and T. G. Sourlingas. Differential sensitivity of human leukemic cell lines to the histone deacetylase inhibitor, trichostatin A. *Leuk. Res.*, 34(6): 786-792, 2009.

-

Happel, N., Doenecke, D., Sekeri-Pataryas, K.E. and T.G. Sourlingas. H1 Histone subtype constitution and phosphorylation state of the ageing cell system of human peripheral blood lymphocytes. *Exp. Gerontol.*, 43: 184-199, 2008.

-

Sourlingas, T.G., Kypreou, K.P., Topakas, G.N. Karchilaki, I.N., Stavropoulou-Giokas, C. and K.E. Sekeri-Pataryas. Effect of the histone deacetylase inhibitor trichostatin A in human peripheral blood lymphocytes as a function of donor age. *Ann. NY Acad. Sci.*, 1119: 64-71, 2007.

-

[Sekeri-Pataryas, K.E. and T.G. Sourlingas.](#) The differentiation-associated linker histone, H1.0, during the in vitro aging and senescence of human diploid fibroblasts. *Ann. N Y Acad. Sci.* 1100: 361-367, 2007.

-

Kypreou, K.P., Sourlingas, T.G. and K.E. Sekeri-Pataryas. Age-dependent response of lymphocytes in the induction of the linker histone variant, H1o and histone H4 acetylation after treatment with the histone deacetylase inhibitor, trichostatin A. *Exp. Gerontol.*, 39(4): 469-479, 2004.

-

Sourlingas, T.G., Issidorides, M.R., Alevizos, B., Kontaxakis, V..P., Chrysanthou-Piterou, M., Livaniou, E., Karpouza, A. and K.E. Sekeri-Pataryas. Lymphocytes from bipolar and schizophrenic patients share common biochemical markers related to histone synthesis and histone cell membrane localization characteristic of an activated state. *Psychiatry Res.*, 118: 55-67, 2003.

-

Sourlingas, T.G., Tsapali, D.S., Kaldis, A.D. and K.E. Sekeri-Pataryas. Histone deacetylase inhibitors induce apoptosis in peripheral blood lymphocytes along with histone H4 acetylation and the expression of the linker histone variant, H1o. *Eur. J. Cell Biol.*, 80: 726-732, 2001.

-

Tsapali. D.S., Sekeri-Pataryas, K.E. and T.G. Sourlingas. mRNA levels of the differentiation-associated linker histone variant, H1 zero in mitotically active and postmitotic senescent human diploid fibroblast cell populations *Exp. Gerontol.*, 36: 1649-1661, 2001.

-

Tsapali, D.S., Sekeri-Pataryas, K.E., Prombona, A. and T.G. Surlingas. mRNA levels of the linker histone variant, H1o, in mitotically active human diploid fibroblasts as a function of the phases of the cell cycle and cumulative population doublings. *Mech. Ageing Dev.* 121 (1-3): 101-112, 2000.

-

Surlingas, T.G., Issidorides, M.R., Havaki, S., Trikkas, G. and K.E. Sekeri-Pataryas. Peripheral blood lymphocytes of bipolar affective patients have a histone synthetic profile indicative of an active cell state. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 22: 81-96, 1998.